

Sharp Regularity near an Absorbing Boundary for Solutions to Second Order SPDEs in a Half-Line with Constant Coefficients

Sean Ledger*

March 28, 2013

Abstract

We prove that the weak version of the SPDE problem

$$\begin{aligned} dV_t(x) &= \left[-\mu V_t'(x) + \frac{1}{2} (\sigma_M^2 + \sigma_I^2) V_t''(x) \right] dt - \sigma_M V_t'(x) dW_t, \quad x > 0 \\ V_t(0) &= 0 \end{aligned}$$

with initial density V_0 a bounded function and W a standard Brownian motion, has a unique solution in the class of finite measure-valued processes. The solution admits a smooth density process which has a probabilistic representation and some degeneracy near the absorbing boundary. We describe the precise order of integrability of the density and its derivatives near the origin via the behaviour of a two-dimensional Brownian motion in a wedge of specific angle. Our results are sharp in the sense that we demonstrate better regularity is unattainable.

1 Introduction

Let W be a standard Brownian motion on a complete filtered probability space $(\Omega, \mathcal{F}, \mathbb{P})$ and V_0 be a deterministic bounded probability density function on the half-line $(0, \infty)$. In this paper we consider the problem of finding a finite-measure valued process $\nu = (\nu_t)_{t \in [0, T]}$ satisfying the weak stochastic partial differential equation

$$\langle \phi, \nu_t \rangle = \langle \phi, V_0 \rangle + \int_0^t \left[\mu \langle \phi', \nu_s \rangle + \frac{1}{2} (\sigma_M^2 + \sigma_I^2) \langle \phi'', \nu_s \rangle \right] ds + \sigma_M \int_0^t \langle \phi', \nu_s \rangle dW_s \quad (1)$$

for all $t \in [0, T]$ and $\phi \in C^{\text{test}}$. We write $\langle \phi, \nu_t \rangle = \int_0^\infty \phi(x) \nu_t(dx)$ and denote the space of bounded, twice-differentiable functions which vanish at the origin and have bounded derivatives by C^{test} . The coefficients $\mu \in \mathbb{R}$, $\sigma_M > 0$ and $\sigma_I > 0$ and the time horizon $T > 0$ are all deterministic constants.

*Mathematical Institute, University of Oxford, 24-29 St Giles, Oxford OX1 3LB, UK. E-mail: sean.ledger@maths.ox.ac.uk

Stochastic partial differential equation (SPDE) problems of this type with zero boundary conditions have been studied extensively [7], [8], [10]. The simplicity of our setting allows greater insight into the behaviour of the solution near the absorbing boundary. We take advantage of a straightforward smoothing technique to present a more elementary study of the problem than is currently offered in the standard literature. We are able to prove uniqueness of solution to (1) in the broad class of finite-measure valued processes. When referring to “uniqueness” throughout this paper, we shall always mean that if ν^1 and ν^2 are two finite-measure valued processes solving (1), then $\mathbb{E} \left[\int_0^T (\langle \phi, \nu_t \rangle - \langle \phi, \bar{\nu}_t \rangle)^2 dt \right] = 0$ for every $\phi \in C^{\text{test}}$.

The results closest to those presented here are found in [8]. In this paper Krylov demonstrates the existence of a unique solution to (1) in the class of twice-differentiable function-valued processes, $u = (u_t(x))_{t \in [0, T]}$, which satisfy

$$\mathbb{E} \left[\int_0^T \left(\|u_t\|_2^2 + \|u'_t\|_2^2 + \|xu''_t\|_2^2 \right) dt \right] < \infty.$$

Furthermore if the initial density is N -times differentiable with $x \mapsto x^k V_0^{(k)} \in L^2(0, \infty)$, for $k = 0, 1, \dots, N$, then it is shown that this solution is $(N + 1)$ -times differentiable with

$$\mathbb{E} \left[\int_0^T \|x^{k-1} u_t^{(k)}\|_2^2 dt \right] < \infty$$

for $k = 0, 1, \dots, N + 1$.

A key ingredient in [8], and the majority of the literature on this problem, is the introduction of weighted Sobolev spaces. These spaces enable one to control the rate at which the solution and its derivatives are allowed to blow-up near the origin. In our one-dimensional setting these weights are easy to visualise and it suffices to use the functions

$$w_c(x) := x^c e^{-x}$$

for $x \in (0, \infty)$ and $c \in \mathbb{R}$. The main result of this paper is to investigate how small one can take c in the above weighting function whilst preserving square-integrability. The critical quantity is the parameter

$$\alpha := \frac{\pi}{2} + \arcsin \frac{\sigma_M}{\sqrt{\sigma_M^2 + \sigma_I^2}} \in \left(\frac{\pi}{2}, \pi \right), \quad (2)$$

which is connected with the underlying dynamics in the probabilistic construction of the solution to (1) presented in section 2. One should observe that the quantity $\pi/\alpha - 1$ appearing in the following theorem is in the range $(0, 1)$.

Theorem 1.1 (The Regularity and Uniqueness Theorem) *Let $N \geq 0$. Suppose that V_0 is bounded and N -times weakly differentiable in $(0, \infty)$ with $w_{-\beta/2} V_0, w_{1-\beta/2} V'_0, w_{2-\beta/2} V''_0, \dots, w_{N-\beta/2} V_0^{(N)}$*

all in $L^2(0, \infty)$ for every $\beta \in (-\infty, \pi/\alpha - 1)$. Then there exists a unique finite measure-valued process, ν , satisfying (1).

Furthermore for almost all $(\omega, t) \in \Omega \times [0, T]$, ν_t has a density V_t that is $(N + 1)$ -times weakly differentiable with

$$\mathbb{E} \left[\int_0^T \left\| w_{k-1-\beta/2} V_t^{(k)} \right\|_2^2 dt \right] < \infty$$

for $k = 0, 1, 2, \dots, N + 1$ and every $\beta \in (-\infty, \pi/\alpha - 1)$.

Theorem 1.1 shows that we gain an extra order of integrability near zero and differentiability in the space dimension as the system evolves forward in time. If one takes $\sigma_M = 0$ then (1) becomes the deterministic heat equation with zero boundary condition. With non-zero σ_M the solution behaves as a heat decay with random fluctuations in location which are driven by the Brownian motion W – see Figure 1. This heat flow behaviour is responsible for the smoothness properties of the density away from the boundary. It is shown in [9], Lemma 3.1 that the density of the solution is infinitely differentiable in the space variable. We can deduce integrability properties of the solution using a comparison with the heat kernel (Proposition 2.5).

As the initial density is bounded and integrable, requiring $w_{k-\beta/2} V_0^{(k)} \in L^2(0, \infty)$ is not a restrictive condition and allows the derivatives of V_0 to be singular at the origin. Notice that since V_0 is bounded and $\beta < 1$, the condition $w_{-\beta/2} V_0 \in L^2(0, \infty)$ is always satisfied. Requiring the condition on V_0 holds for arbitrarily negative values of β does not substantially reduce the class of admissible initial densities, since the weighting functions decay exponentially at infinity. We therefore allow initial densities whose derivatives grow sub-exponentially in magnitude towards infinity. In Theorem 1.1 the case $\beta = 0$ corresponds to the aforementioned result in [8], and hence our results expand upon the known regularity of the solution at the boundary.

Naturally one would like to know whether we can take $\beta > \pi/\alpha - 1$ in Theorem 1.1. It is shown in [6], Lemma 5.1 that the critical estimate in Theorem 2.6 required in the proof of Theorem 1.1 is sharp, and in turn this allows us to show that the regularity is also sharp:

Theorem 1.2 (Converse result) *Assume that $V_0 \in L^2(0, \infty)$ is bounded and N -times weakly differentiable in $(0, \infty)$ with $w_{-\beta'/2} V_0, w_{1-\beta'/2} V_0', w_{2-\beta'/2} V_0'', \dots, w_{N-\beta'/2} V_0^{(N)}$ all in $L^2(0, \infty)$ for every $\beta' \in (-\infty, \pi/\alpha - 1)$. (One can allow V_0 to be smoother by extending this range of β' to $\mathbb{R}$ and the conclusion is unchanged.) Then for every $\beta \in (\pi/\alpha - 1, \infty)$*

$$\mathbb{E} \left[\int_0^T \left\| w_{k-1-\beta/2} V_t^{(k)} \right\|_2^2 dt \right] = \infty$$

for $k = 0, 1, \dots, N$.

The blow-up of the second derivative is an interesting feature of SPDE problems that one does not observe in the deterministic setting – that is when $\sigma_M = 0$. In [8], Example 1.2 a system similar

to (1) is presented for which the solution does not have second derivative that is in $L^2(0, \infty)$. An explicit description of the blow-up is given in [9]. Suppose $\mu = 0$ and shift the density of the solution by setting

$$f(t, x) := V_t(x + \sigma_M W_t).$$

Then f solves the heat equation (with drift) in the region

$$\{(t, x) : t \in [0, T], x > -\sigma_M W_t\}$$

with zero boundary condition. It is shown (Theorem 5.1, [9]) that there exists $\lambda \in (0, 1)$ such that, with probability one, there is a dense subset of $[0, T]$ on which

$$\lim_{x \downarrow 0} \frac{f(t, x)}{x^\lambda} = \infty.$$

On this set it is clear that the density can have no space derivative at the absorbing boundary.

Our study of this SPDE problem is motivated by [4], in which a structural model is proposed for the loss due to default in a large portfolio of credit risky assets. The authors construct a finite system of Brownian motions, correlated through a common diffusion term, which are killed upon hitting a boundary at zero. As the number of particles in the system approaches infinity, the empirical measure of the particles – that is the proportion of particles falling in a given set – converges to the probabilistic solution ν of (1) defined in Section 2. That (1) has a unique measure-valued solution allows us to characterise the model in [4] through the SPDE and Dirichlet boundary condition.

In the proceeding section we present the construction of the solution to (1) and demonstrate its essential properties. A proof of the significant $(3 + \beta)$ -condition (Theorem 2.6) is presented in Section 3. The remaining sections are devoted to proving the above two theorems.

2 The Probabilistic Solution

2.1 Construction

Let us introduce another complete probability space $(\Omega^I, \mathcal{F}^I, \mathbb{P}^I)$ and independent standard Ω^I -Brownian motions W^1 and W^2 . We will denote expectation under $\mathbb{P}^I$ by $\mathbb{E}^I$. Let the processes $\tilde{X}^1$ and $\tilde{X}^2$ be defined as

$$\tilde{X}_t^i := X_0^i + \mu t + \sigma_M W_t + \sigma_I W_t^i$$

where X_0^1, X_0^2 are independent Ω^I -random variables with common law V_0 . We stop the particles upon hitting zero by setting

$$X_t^i := \begin{cases} \tilde{X}_t^i & t \in [0, \tau_i) \\ \partial & t \in [\tau_i, T] \end{cases}$$




Figure 1: Presented on the left is the evolution of the system in the deterministic case $\sigma_M = 0$. The value of the density at a space-time point is represented by a colour increasing from blue ($V_t(x) = 0$) through to red ($V_t(x) = 1$). The initial condition is a step function. Compare this with a realisation of the evolution for some $\sigma_M > 0$, which is presented on the right. We observe heat-type dispersion but with a non-smooth fluctuation.

where ∂ is some cemetery state (one can take $\partial = 0$ for our purposes) and

$$\tau_i := \inf \left\{ s \in [0, T] : \tilde{X}_s^i \leq 0 \right\}.$$

Clearly X^1 and X^2 are $\Omega \times \Omega^I$ -random variables, and conditional on knowing $\omega \in \Omega$, $X^1(\omega, \cdot)$ and $X^2(\omega, \cdot)$ are independent Ω^I -random variables. It is therefore immediate that X^1 and X^2 are equal in Ω^I -distribution, so we make the following definition:

Definition 2.1 (Probabilistic Solution) *Let the probabilistic solution, ν , be the measure-valued process defined by*

$$\nu_t(S) := \mathbb{P}^I(X_t^1 \in S)$$

for all $t \in [0, T]$ and $S \subseteq (0, \infty)$ Lebesgue measurable. Notice that ν is a Ω -random variable.

Throughout this paper “ $S \subseteq (0, \infty)$ measurable” shall always refer to a Lebesgue measurable subset S . The following useful:

Proposition 2.2 *For all $t \in [0, T]$ and $S \subseteq (0, \infty)$ measurable we have*

$$\mathbb{E} \left[\nu_t(S)^2 \right] = \mathbb{E} \left[\mathbb{E}^I \left[\mathbf{1}_{X_t^1 \in S, X_t^2 \in S} \right] \right]$$

Proof: Since X^1 and X^2 are equal in Ω^I -distribution we have $\nu_t(S) = \mathbb{E}^I [\mathbf{1}_{X_t^i \in S}]$ for $i = 1, 2$. Hence

$$\mathbb{E} [\nu_t(S)^2] = \mathbb{E} \left[\mathbb{E}^I [\mathbf{1}_{X_t^1 \in S}]^2 \right] = \mathbb{E} \left[\mathbb{E}^I [\mathbf{1}_{X_t^1 \in S}] \mathbb{E}^I [\mathbf{1}_{X_t^2 \in S}] \right] = \mathbb{E} \left[\mathbb{E}^I [\mathbf{1}_{X_t^1 \in S, X_t^2 \in S}] \right]$$

as required, where the final line is due to the Ω^I -independence of X^1 and X^2 . □

It is a matter of applying Ito's formula to see that ν is indeed a solution of (1):

Proposition 2.3 *There exists a full subset of Ω on which*

$$\langle \phi, \nu_t \rangle = \langle \phi, V_0 \rangle + \int_0^t \left[\mu \langle \phi', \nu_s \rangle + \frac{1}{2} (\sigma_M^2 + \sigma_I^2) \langle \phi'', \nu_s \rangle \right] ds + \sigma_M \int_0^t \langle \phi', \nu_s \rangle dW_s.$$

for all $t \in [0, T]$ and $\phi \in C^{test}$.

Proof: We need to be careful near the boundary at zero. Begin by extending ϕ onto $\mathbb{R}$, for example by setting $\phi(x) := \frac{1}{2}\phi''(0)x^2 + \phi'(0)x$ for $x < 0$. Then $\phi \in C^2(\mathbb{R})$ so we can apply Ito's formula to $\phi(\tilde{X})$ which gives that for almost all $\omega \in \Omega$

$$\begin{aligned} \phi(\tilde{X}_t^1) &= \phi(X_0^1) + \int_0^t \left[\mu \phi'(\tilde{X}_s^1) + \frac{1}{2} (\sigma_M^2 + \sigma_I^2) \phi''(\tilde{X}_s^1) \right] ds + \sigma_M \int_0^t \phi'(\tilde{X}_s^1) dW_s \\ &\quad + \sigma_I \int_0^t \phi'(\tilde{X}_s^1) dW_s^1 \end{aligned}$$

for all $t \in [0, T]$. We can replace t by $t \wedge \tau$, noting that $\phi(\tilde{X}_{t \wedge \tau}^1) = \phi(X_{t \wedge \tau}^1)$, since $\phi(0) = \phi(\tilde{X}_\tau^1) = 0 = \phi(\partial)$, and

$$\int_0^t \phi'(X_s^1) ds = \int_0^{t \wedge \tau} \phi'(X_s^1) ds + \int_{t \wedge \tau}^t \phi'(\partial) ds = \int_0^{t \wedge \tau} \phi'(\tilde{X}_s^1) ds$$

and likewise for the other integral terms, therefore

$$d\phi(X_t^1) = \left[\mu \phi'(X_t^1) + \frac{1}{2} (\sigma_M^2 + \sigma_I^2) \phi''(X_t^1) \right] dt + \sigma_M \phi'(X_t^1) dW_t + \sigma_I \phi'(X_t^1) dW_t^1$$

and the result follows by taking expectation over Ω^I and noting that $\mathbb{E}^I [\phi(X_t^1)] = \langle \phi, \nu_t \rangle$. □

Remark 2.4 (Connection with filtering) If we treat X^1 as a signal process and W an observation process, then the Zakai equation (see for example [1]) for the conditional distribution of the signal given the observations is exactly (1).

2.2 Properties

For a fixed $\omega \in \Omega$ the process $\tilde{X}^1$ which is not killed upon hitting the origin is simply a Ω^I -Brownian motion started at X_0^1 with drift $t \mapsto \mu t + \sigma_M W_t$. We can therefore explicitly write the Ω^I -density of $\tilde{X}^1$ as:

$$P_t^I(x; x_0) = P_t^I(x; x_0)(\omega) = \frac{1}{\sqrt{2\pi\sigma_I^2 t}} \exp \left\{ -\frac{(x - x_0 - \mu t - \sigma_M W_t)^2}{2\sigma_I^2 t} \right\}.$$

It is trivial to see that if $X_t^1 \in S$ for a time $t \in [0, T]$ and a measurable subset $S \subseteq (0, \infty)$, then one must have $\tilde{X}_t^1 \in S$. This simple observation allows us to show that ν has a density process:

Proposition 2.5 (Existence of the density) *There exists a full subset of Ω on which for every $t \in [0, T]$, ν_t has a density process $V_t \in L^p(0, \infty)$ for all $1 \leq p \leq \infty$, with $\|V_t\|_p \leq \|V_0\|_p$.*

Proof: From the above observation, there exists a full subset Ω on which

$$\nu_t(S) \leq \mathbb{P}^I(\tilde{X}_t^1 \in S) = \int_S \int_0^\infty P_t^I(x; x_0) V_0(dx_0) dx$$

for all $t \in [0, T]$ and $S \subseteq (0, \infty)$ measurable. Therefore if $S \subseteq (0, \infty)$ has zero Lebesgue measure then $\nu_t(S) = 0$. It follows by the Radon–Nikodym Theorem that ν_t has a density for every $t \in [0, T]$, and that this density V_t satisfies

$$V_t(x) \leq \int_0^\infty P_t^I(x; x_0) V_0(x_0) dx_0 \quad (3)$$

for almost all $x \in (0, \infty)$.

Bounding V_0 by $\|V_0\|_\infty$ in (3), we get $V_t \in L^\infty(0, \infty)$ with $\|V_t\|_\infty \leq \|V_0\|_\infty$. Suppose $1 \leq p < \infty$, then Holder's inequality applied to (3) gives

$$V_t(x)^p \leq \int_0^\infty P_t^I(x; x_0) V_0(x_0)^p dx_0.$$

Integrating over $x \in (0, \infty)$ and noting that $\int_0^\infty P_t^I(x; x_0) dx \leq 1$ gives $\|V_t\|_p \leq \|V_0\|_p$.

□

The most important feature of the probabilistic solution is the following estimate:

Theorem 2.6 (The $(3 + \beta)$ -Condition) *Let $V_0 \in L^2(0, \infty)$ be bounded and α be defined as in (2). Then for every $\beta \in (0, \pi/\alpha - 1) \subseteq (0, 1)$ there exists constants $B > 0$ and $\gamma \in (0, 1)$ such that for all $t \in [0, T]$ and $\varepsilon > 0$*

$$\mathbb{E} \left[\nu_t(0, \varepsilon)^2 \right] \leq \frac{B}{t^\gamma} \varepsilon^{3+\beta}.$$

A proof is presented in section 3.

In Lemma 3.5 of [4] this result is presented for an initial density supported on a finite interval away from the boundary. This assumption allows the authors to drop the singular $t^{-\gamma}$ factor. However in some respects this is not a natural initial condition. For any $t > 0$, V_t takes positive values on all of $(0, \infty)$, so if we stop the process at time t_0 and then restart from the initial density V_{t_0} we still have the estimate on $\mathbb{E} \left[\nu_t(0, \varepsilon)^2 \right]$ for $t > t_0$, despite V_{t_0} not being supported on a finite interval away from zero. In this more general setting we can expect a singular time factor to appear since for every time $t > 0$, $V_t(x) \rightarrow 0$ as $x \rightarrow 0$ (Theorem 3.2 of [9]). Therefore positive values of V_0 close to zero must decay instantaneously as the system evolves in time. Since $\gamma \in (0, 1)$, $t^{-\gamma}$ is integrable over $[0, T]$ and so this factor does not present technical difficulties in later proofs involving L^2 -integrability.

Using Proposition 2.2 we prove Theorem 2.6 by considering the process $\mathbf{X}_t = (X_t^1, X_t^2)$. Under $\mathbb{P} \times \mathbb{P}^I$ this process is a two-dimensional Brownian motion that is killed upon exiting the quadrant $(0, \infty)^2$ and has components with correlation

$$\rho = \frac{\sigma_M}{\sqrt{\sigma_M^2 + \sigma_I^2}}.$$

With α as in (2), the map $F_\alpha : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by

$$F_\alpha : \mathbf{x} \mapsto \begin{pmatrix} \sqrt{1-\rho^2} & \rho \\ 0 & 1 \end{pmatrix}^{-1} \mathbf{x} \quad (4)$$

transforms $\mathbf{X}$ to a Brownian motion with uncorrelated components that is killed upon exiting the wedge

$$F_\alpha \left[(0, \infty)^2 \right] = \{(r, \theta) : 0 < r < \infty, 0 < \theta < \alpha\}.$$

We then make use of explicit formulae in [5], [12] to estimate the hitting probability of a small neighborhood of the apex of the wedge. The kernel smoothing technique introduced in [4] and adapted from [11] then allows us to relate this hitting probability to the behaviour of the solution near the absorbing boundary.

3 Proof of Theorem 2.6

Let us write $\mathbb{P}_{x_1, x_2}$ to indicate that $X_0^1 = x_1$ and $X_0^2 = x_2$, that is we start the two-dimensional Brownian motion $\mathbf{X} = (X^1, X^2)$ from (x_1, x_2) . From Proposition 2.2 we know

$$\mathbb{E}_{x_1, x_2} [\nu_t(S_1) \nu_t(S_2)] = \mathbb{E}_{x_1, x_2} \left[\mathbb{E}^I \left[\mathbf{1}_{X_t^1 \in S_1, X_t^2 \in S_2} \right] \right] \quad (5)$$

for all $S_1, S_2 \subseteq (0, \infty)$ measurable. A change of measure allows us to eliminate the drift term in the dynamics of $\mathbf{X}$:

Lemma 3.1 (Eliminating drift) *Let $\tilde{\mathbb{P}}$ be defined on $(\Omega, \mathcal{F}_t)$ by the Girsanov transformation*

$$\frac{d\tilde{\mathbb{P}}}{d\mathbb{P}} = Z_t(\mu) := \exp \left\{ -\frac{\mu}{\sigma_M} W_t - \frac{\mu^2}{2\sigma_M^2} t \right\}$$

so that under $\tilde{\mathbb{P}} \times \mathbb{P}^I$ $\mathbf{X}$ is a two-dimensional $(\Omega \times \Omega^I)$ -Brownian motion with zero drift. Let $\tilde{\mathbb{E}}$ denote the expectation under $\tilde{\mathbb{P}}$. If there exists a constant $C > 0$ such that

$$\tilde{\mathbb{E}} \left[\nu_t(0, \varepsilon)^2 \right] \leq \frac{C}{t^{\pi/2\alpha}} \varepsilon^{2+\pi/\alpha}$$

for all $t \in (0, T]$ and $\varepsilon > 0$, then Theorem 2.6 holds.

Proof: Following the method in Lemma 3.5 of [4], for any Holder conjugates $1 < a, b < \infty$ we have

$$\begin{aligned} \mathbb{E} \left[\nu_t(0, \varepsilon)^2 \right] &= \tilde{\mathbb{E}} \left[Z_t(\mu) \cdot \nu_t(0, \varepsilon)^2 \right] \\ &\leq \tilde{\mathbb{E}} \left[Z_t(\mu)^a \right]^{\frac{1}{a}} \tilde{\mathbb{E}} \left[\nu_t(0, \varepsilon)^{2b} \right]^{\frac{1}{b}} \\ &\leq \tilde{\mathbb{E}} \left[Z_t(\mu)^a \right]^{\frac{1}{a}} \tilde{\mathbb{E}} \left[\nu_t(0, \varepsilon)^2 \right]^{\frac{1}{b}} \\ &\leq \tilde{\mathbb{E}} \left[Z_t(\mu)^a \right]^{\frac{1}{a}} \left\{ \frac{C}{t^{\pi/2\alpha}} \varepsilon^{2+\pi/\alpha} \right\}^{\frac{1}{b}} \\ &= \mathbb{E} \left[Z_t(\mu)^{a-1} \right]^{\frac{1}{a}} \left\{ \frac{C}{t^{\pi/2\alpha}} \varepsilon^{2+\pi/\alpha} \right\}^{\frac{1}{b}} \end{aligned}$$

where one should note that $\nu_t(0, \varepsilon) \leq 1$. Hence

$$\mathbb{E} \left[\nu_t(0, \varepsilon)^2 \right] \leq \exp \left(\frac{(a-1)(a-2)\mu^2}{2\sigma_M^2 a} T \right) \left\{ \frac{C}{t^{\pi/2\alpha}} \varepsilon^{2+\pi/\alpha} \right\}^{\frac{1}{b}},$$

therefore we can choose b sufficiently close to 1 so that the result holds since $\alpha \in (\pi/2, \pi)$.

□

Remark 3.2 Notice that we cannot take $b = 1$ above. This change of measure technique eliminates the possibility of taking $\beta = \pi/\alpha - 1$, except in the case when $\mu = 0$. Although this may seem to weaken the potential regularity available near the default boundary, in proving Theorem 1.1 we make use of Lemma 4.8 which requires β to be strictly smaller than this maximal value. It is therefore unclear as to the precise behaviour of V at the critical value $\beta = \pi/\alpha - 1$.

The remainder of Section 3 is devoted to showing the hypothesis of Lemma 3.1 holds. Clearly it is sufficient to take $\mu = 0$ and $\tilde{\mathbb{P}} = \mathbb{P}$.

Conditioning on the start-point of $\mathbf{X}$:

$$\mathbb{E} \left[\nu_t(0, \varepsilon)^2 \right] = \int_0^\infty \int_0^\infty \mathbb{E}_{x_1, x_2} \left[\mathbb{E}^I \left[\mathbf{1}_{X_t^1 \in (0, \varepsilon), X_t^2 \in (0, \varepsilon)} \right] \right] V_0(x_1) V_0(x_2) dx_1 dx_2.$$

Now the region $(0, \varepsilon)^2$ is mapped *into* a region of the form $\{(r, \theta) : 0 < r < c\varepsilon, 0 < \theta < \alpha\}$, for some numerical constant c , under the transformation F_α . As the estimate we are demonstrating is unchanged under the map $\varepsilon \mapsto c\varepsilon$ (modulo multiplicative constants), it is no loss of generality to take $c = 1$. Using the fact that V_0 is bounded, that the Jacobian of the transform $F_\alpha : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a numerical constant depending only on ρ , and formula (8) of [5] (corroborated in [12]), we arrive at

$$\begin{aligned} \mathbb{E} \left[\nu_t(0, \varepsilon)^2 \right] &\leq C \int_0^\infty \int_0^\alpha \int_0^\alpha \int_0^\varepsilon \frac{rr_0}{t} e^{-\frac{r^2+r_0^2}{2t}} \sum_{n=1}^\infty \sin \frac{n\pi\theta}{\alpha} \sin \frac{n\pi\theta_0}{\alpha} I_{n\pi/\alpha} \left(\frac{rr_0}{2t} \right) dr d\theta dr_0 d\theta_0 \\ &= C \int_0^\varepsilon \frac{r^{\pi/\alpha+1}}{t^{\pi/2\alpha}} e^{-\frac{r^2}{2t}} \sum_{n,m=0}^\infty \frac{\Gamma \left(m + \frac{(2n+1)\pi}{2\alpha} + 1 \right)}{(2n+1)^2 m! \Gamma \left(m + \frac{(2n+1)\pi}{\alpha} + 1 \right)} \left(\frac{r}{2t} \right)^{m+n\pi/\alpha} dr \end{aligned}$$

for $C > 0$ a numerical constant. As $\sup_{x \geq 0} \{x^c e^{-x}\} = c^c e^{-c}$ we have

$$\mathbb{E} \left[\nu_t(0, \varepsilon)^2 \right] \leq C \int_0^\varepsilon \frac{r^{\pi/\alpha+1}}{t^{\pi/2\alpha}} dr \sum_{n,m=0}^\infty \frac{\Gamma \left(m + \frac{(2n+1)\pi}{2\alpha} + 1 \right) \left(m + \frac{n\pi}{\alpha} \right)^{m+\frac{n\pi}{\alpha}} e^{-(m+\frac{n\pi}{\alpha})}}{(2n+1)^2 m! \Gamma \left(m + \frac{(2n+1)\pi}{\alpha} + 1 \right)}. \quad (6)$$

So we are finished provided the double sum on the right-hand is finite. Using Stirling's approximation, the inequality

$$\frac{\Gamma(a+c+1)}{\Gamma(b+c+1)} \leq \frac{\Gamma(a+1)}{\Gamma(b+1)}$$

for $a, b, c > 0$ and the asymptotic behaviour $\Gamma(m + \pi/2\alpha + 1) \sim m! m^{\pi/2\alpha}$ for large m , gives that the summand in (6) is dominated by a constant multiple of

$$\frac{1}{(2n+1)^2 m^{\pi/2\alpha+1/2}}$$

for large m and $m+n$. Since $\pi/2\alpha + 1/2 > 1$, this expression is summable over m and n and so we have the result. □

4 Proof of Theorem 1.1

We begin by proving the regularity result for the probabilistic solution. Let us write p_δ for the transition density of a Brownian motion in $\mathbb{R}$ with zero drift and unit volatility, that is

$$p_\delta(x) = \frac{1}{\sqrt{2\pi\delta}} \exp\left\{-\frac{x^2}{2\delta}\right\} \quad (7)$$

for $z \in \mathbb{R}$, $\delta > 0$.

Definition 4.1 (Heat kernels) *Let $\delta > 0$. The absorbing and reflected heat kernels are defined to be*

$$\begin{aligned} G_\delta(x, y) &:= p_\delta(x - y) - p_\delta(x + y) \\ G_\delta^r(x, y) &:= p_\delta(x - y) + p_\delta(x + y) \end{aligned}$$

for $x, y > 0$, respectively. The smoothed density and reflected smoothed density are then be defined to be

$$\begin{aligned} T_{t,\delta}(x) &:= \int_0^\infty G_\delta(x, y) V_t(y) dy \\ T_{t,\delta}^r(x) &:= \int_0^\infty G_\delta^r(x, y) V_t(y) dy \end{aligned}$$

for $t \in [0, T]$ and $x > 0$, respectively.

Now take any smooth compactly supported function $\phi \in C_K^\infty(0, \infty)$. Then for every $\delta > 0$, $x \mapsto T_\delta\phi(x)$ (that is the function obtained by convolving the heat kernel with ϕ) is in C^{test} , so putting $T_\delta\phi$ and ν into (1) we arrive at the following strong equation for the evolution of the smoothed density:

Proposition 4.2 (Strong smoothed evolution equation) *For $\delta > 0$, $t \in [0, T]$ and $x > 0$*

$$\begin{aligned} T_{t,\delta}(x) &= T_{0,\delta}(x) + \int_0^t \left[-\mu T'_{s,\delta}(x) + \frac{\sigma_M^2 + \sigma_I^2}{2} T''_{s,\delta}(x) \right] ds - \sigma_M \int_0^t T'_{s,\delta}(x) dW_s \\ &\quad + \mu \int_0^t R'_{s,\delta}(x) ds + \sigma_M \int_0^t R'_{s,\delta}(x) dW_s. \end{aligned} \quad (8)$$

where $R_{t,\delta}$ is the remainder function:

$$R_{t,\delta}(x) := T_{t,\delta}(x) - T_{t,\delta}^r(x) = 2 \int_0^\infty p_\delta(x + y) V_t(y) dy.$$

Proof: This is simply a matter of observing that

$$\begin{aligned}\langle \phi', T_{t,\delta} \rangle &= -\langle \phi, (T_{t,\delta}^r)' \rangle \\ \langle \phi'', T_{t,\delta} \rangle &= \langle \phi, T_{t,\delta}'' \rangle.\end{aligned}$$

One also needs to apply the stochastic Fubini theorem [3] to switch integration in the space dimension with integration with respect to W (and likewise with respect to time). To this end it is enough to note that the tails of $T_{t,\delta}$ and $T_{t,\delta}^r$ and their derivatives decay exponentially, hence the conditions of the (stochastic) Fubini theorem are indeed met.

□

Differentiating equation (8) n times (or integrating once for the case $n = -1$, note the result is no different) and applying Ito's formula to $\left(T_{t,\delta}^{(n)}\right)^2$ gives:

$$\begin{aligned}d\left(T_{t,\delta}^{(n)}\right)^2 &= -2\mu T_{t,\delta}^{(n)} T_{t,\delta}^{(n+1)} dt + (\sigma_M^2 + \sigma_I^2) T_{t,\delta}^{(n)} T_{t,\delta}^{(n+2)} dt + \sigma_M^2 \left(T_{t,\delta}^{(n+1)}\right)^2 dt \\ &\quad + \sigma_M^2 \left(R_{t,\delta}^{(n+1)}\right)^2 dt + 2\mu T_{t,\delta}^{(n)} R_{t,\delta}^{(n+1)} dt \\ &\quad - 2\sigma_M T_{t,\delta}^{(n)} T_{t,\delta}^{(n+1)} dW_t + 2\sigma_M T_{t,\delta}^{(n)} R_{t,\delta}^{(n+1)} dW_t.\end{aligned}\tag{9}$$

From this equation we shall proceed to the proof by induction in section 4.2. Before we continue we introduce several technical lemmas.

4.1 Technical Lemmas

Lemma 4.3 *There exists a full subset of Ω on which $x^{-c}T_{t,\delta}(x) \rightarrow 0$ as $x \rightarrow 0$ for every $c \in (0, 1)$, $\delta > 0$ and $t \in (0, T]$.*

Proof: This is straightforward since for $x < 1$ we have

$$\begin{aligned}0 \leq T_{t,\delta}(x) &= \frac{1}{\sqrt{2\pi\delta}} \int_0^\infty \left[\exp\left\{-\frac{(x-y)^2}{2\delta}\right\} - \exp\left\{-\frac{(x+y)^2}{2\delta}\right\} \right] V_t(y) dy \\ &\leq \frac{2\|V_0\|_\infty x}{\delta\sqrt{2\pi\delta}} \int_0^\infty y \exp\left\{-\frac{(y-1)^2}{2\delta}\right\} dy \\ &= D_\delta x\end{aligned}$$

for a numerical constant $D_\delta > 0$, and where we have used the estimate $1 - e^{-z} \leq z$.

□

Lemma 4.4 *Let $n \geq 0$ be an integer. Suppose that for $k = 0, 1, \dots, n$*

$$\liminf_{\delta \rightarrow 0} \mathbb{E} \left[\int_0^T \left\| w_{k-1-\beta/2} (T_{t,\delta})^{(k)} \right\|_2^2 dt \right] < \infty.$$

Then for almost all $(\omega, t) \in \Omega \times [0, T]$, the density V_t is n -times weakly differentiable with

$$\mathbb{E} \left[\int_0^T \left\| w_{k-1-\beta/2} V_t^{(k)} \right\|_2^2 dt \right] < \infty \quad (10)$$

for $k = 0, 1, \dots, n$.

Proof: Let $\langle \cdot, \cdot \rangle$ denote the usual $L^2(0, \infty)$ inner product: $\langle f, g \rangle = \int_0^\infty f(x) g(x) dx$. Let $\{\phi_i\}_{i=1}^\infty$ be a basis of $L^2(0, \infty)$ such that each ϕ_i is smooth and bounded.

Firstly for any $\phi \in C^\infty(0, \infty)$ and $m > k \geq 1$, $\lim_{x \downarrow 0} (w_{m-\beta/2} \phi(x))^{(k)} = 0$, since $\partial_x^k w_{m-\beta/2}$ is a linear combination of $w_{i-\beta/2} \rightarrow 0$, for $m-k \leq i \leq m$. So for $k \geq 1$, integrating by parts gives

$$\left\langle (w_{k-1-\beta/2} \phi)^{(k)}, T_{t,\delta} \right\rangle = (-1)^k \left\langle \phi, w_{k-1-\beta/2} T_{t,\delta}^{(k)} \right\rangle - \lim_{x \downarrow 0} \left\{ T_{t,\delta}(x) (w_{k-1-\beta/2} \phi)^{(k-1)} \right\}.$$

Applying Leibniz's rule to $(w_{k-1-\beta/2} \phi)^{(k-1)}$, we see that

$$\lim_{x \downarrow 0} \left\{ T_{t,\delta}(x) (w_{k-1-\beta/2} \phi)^{(k-1)}(x) \right\} = C_k \lim_{x \downarrow 0} \left\{ T_{t,\delta}(x) w_{-\beta/2}(x) \phi(x) \right\}$$

for some numerical constant C_k , and the right-hand-side vanishes by Lemma 4.3, since $w_{-\beta/2}(x) = \mathcal{O}(x^{-\beta/2})$ for small x . Hence we conclude

$$\left\langle (w_{k-1-\beta/2} \phi)^{(k)}, T_{t,\delta} \right\rangle = (-1)^k \left\langle \phi, w_{k-1-\beta/2} T_{t,\delta}^{(k)} \right\rangle.$$

The remaining case $k = 0$ is trivial.

From these results and Fatou's Lemma we have

$$\begin{aligned} \mathbb{E} \left[\int_0^T \sum_{i=1}^\infty \left| \left\langle (w_{k-1-\beta/2} \phi_i)^{(k)}, V_t \right\rangle \right|^2 dt \right] &= \mathbb{E} \left[\int_0^T \sum_{i=1}^\infty \lim_{\delta \rightarrow 0} \left| \left\langle (w_{k-1-\beta/2} \phi_i)^{(k)}, T_{t,\delta} \right\rangle \right|^2 dt \right] \\ &\leq \liminf_{\delta \rightarrow 0} \mathbb{E} \left[\int_0^T \sum_{i=1}^\infty \left| \left\langle \phi_i, w_{k-1-\beta/2} T_{t,\delta}^{(k)} \right\rangle \right|^2 dt \right] \\ &= \liminf_{\delta \rightarrow 0} \mathbb{E} \left[\int_0^T \left\| w_{k-1-\beta/2} T_{t,\delta}^{(k)} \right\|_2^2 dt \right] \\ &< \infty \end{aligned}$$

for $k = 0, 1, \dots, n$, therefore we can define the processes

$$W_{t,k}(x) := \sum_{i=1}^{\infty} \left\langle (w_{k-1-\beta/2} \phi_i)^{(k)}, V_t \right\rangle \phi_i(x)$$

which satisfy

$$\mathbb{E} \left[\int_0^T \|W_{t,k}\|_2^2 dt \right] < \infty.$$

For $k = 0$, let $f_0 = w_{-1-\beta/2}$ and take any $\psi \in C_K^\infty(0, \infty)$, then we calculate

$$\left\langle \psi, \frac{1}{f_0} W_{t,0} \right\rangle = \sum_{i=1}^{\infty} \langle f_0 \phi_i, V_t \rangle \left\langle \frac{\psi}{f_0}, \phi_i \right\rangle = \left\langle f_0 \sum_{i=1}^{\infty} \left\langle \frac{\psi}{f_0}, \phi_i \right\rangle \phi_i, V_t \right\rangle = \langle \psi, V_t \rangle$$

and so $V_t = W_{t,0}/f_0$. Hence we have demonstrated (10) for $k = 0$.

Take $k > 0$, and set $f_k = w_{k-1-\beta/2}$. Using $V_t = W_{t,0}/f_0$ we can compute

$$\begin{aligned} \left\langle \psi, \frac{1}{f_k} W_{t,k} \right\rangle &= \sum_{i=1}^{\infty} \left\langle (f_k \phi_i)^{(k)}, V_t \right\rangle \left\langle \frac{\psi}{f_k}, \phi_i \right\rangle \\ &= \sum_{i,j=1}^{\infty} \left\langle (f_k \phi_i)^{(k)}, \frac{\phi_j}{f_0} \right\rangle \langle f_0 \phi_j, V_t \rangle \left\langle \frac{\psi}{f_k}, \phi_i \right\rangle \\ &= \sum_{i,j,m=1}^{\infty} \left\langle (f_k \phi_i)^{(k)}, \phi_m \right\rangle \left\langle \frac{\phi_j}{f_0}, \phi_m \right\rangle \langle f_0 \phi_j, V_t \rangle \left\langle \frac{\psi}{f_k}, \phi_i \right\rangle \\ &= \sum_{i,j,m=1}^{\infty} (-1)^k \left\langle \phi_i, f_k \phi_m^{(k)} \right\rangle \left\langle \frac{\phi_j}{f_0}, \phi_m \right\rangle \langle f_0 \phi_j, V_t \rangle \left\langle \frac{\psi}{f_k}, \phi_i \right\rangle \\ &= \sum_{j,m=1}^{\infty} (-1)^k \left\langle \psi, \phi_m^{(k)} \right\rangle \left\langle \frac{\phi_j}{f_0}, \phi_m \right\rangle \langle f_0 \phi_j, V_t \rangle \\ &= \sum_{j=1}^{\infty} \left\langle \psi^{(k)}, \frac{\phi_j}{f_0} \right\rangle \langle f_0 \phi_j, V_t \rangle \\ &= \left\langle \psi^{(k)}, V_t \right\rangle, \end{aligned}$$

hence $(-1)^k W_{t,k}/f_k$ is the k^{th} weak derivative of V_t , which completes the proof. □

Lemma 4.5 *There exists a full subset of Ω on which*

$$\int_0^\infty w_{n-\beta/2}^2 T_{t,\delta}^{(n)}(x) T_{t,\delta}^{(n+1)}(x) dx = \left\| w_{n-\beta/2} T_{t,\delta}^{(n)} \right\|_2^2 - \left(n - \frac{\beta}{2} \right) \left\| w_{n-1/2-\beta/2} T_{t,\delta}^{(n)} \right\|_2^2$$

and

$$\begin{aligned}
\int_0^\infty w_{n-\beta/2}^2 T_{t,\delta}^{(n)}(x) T_{t,\delta}^{(n+2)}(x) dx &= 2 \left\| w_{n-\beta/2} T_{t,\delta}^{(n)} \right\|_2^2 - \left\| w_{n-\beta/2} T_{t,\delta}^{(n+1)} \right\|_2^2 \\
&\quad - 4 \left(n - \frac{\beta}{2} \right) \left\| w_{n-1/2-\beta/2} T_{t,\delta}^{(n)} \right\|_2^2 \\
&\quad + 2 \left(n - \frac{\beta}{2} \right) \left(n - \frac{\beta+1}{2} \right) \left\| w_{n-1-\beta/2} T_{t,\delta}^{(n)} \right\|_2^2
\end{aligned}$$

for every $n \geq -1$, $\delta > 0$ and $t \in [0, T]$.

Proof: The result is an integration-by-parts exercise provided the following limits vanish:

$$\begin{aligned}
&\lim_{x \downarrow 0} \left\{ w_{n-1/2-\beta/2}(x) T_{t,\delta}^{(n)}(x) \right\} \\
&\lim_{x \downarrow 0} \left\{ w_{n-\beta/2}^2(x) T_{t,\delta}^{(n)}(x) T_{t,\delta}^{(n+1)}(x) \right\}
\end{aligned}$$

for $n \geq -1$. If $n \geq 1$ then $w_{n-1/2-\beta/2}, w_{n-\beta/2} \rightarrow 0$ as $x \rightarrow 0$, and the result is immediate. In the case $n = 0$, Lemma 4.3 gives $T_{t,\delta}(x) = \mathcal{O}(x)$ which is sufficient for the limit to be zero. This also implies $T_{t,\delta}^{(-1)}(x) = \mathcal{O}(x^2)$, so the result holds for $n = -1$ too.

□

Lemma 4.6 With p_δ defined as in (7), for every integer $n \geq 0$ there exist constants c_i^n for $n/2 \leq i \leq n$ such that for every $\delta > 0$ and $x > 0$

$$\frac{\partial^n p_\delta}{\partial x^n}(x) = \sum_{n/2 \leq i \leq n} c_i^n x^{2i-n} \delta^{-i} p_\delta(x).$$

Proof: This is a simple inductive argument.

□

Lemma 4.7 For $n = -1, 0, 1, \dots, N$ and $\beta < \pi/\alpha - 1$

$$\liminf_{\delta \rightarrow 0} \left\| w_{n-\beta/2} T_{0,\delta}^{(n)} \right\|_2^2 < \infty.$$

Proof: The cases $n = 0$ and $n = -1$ are simple. Firstly observe from Definition 4.1 that $\|T_{0,\delta}\|_\infty \leq \|V_0\|_\infty$. Therefore for all $\delta > 0$

$$\left\| w_{-\beta/2} T_{0,\delta} \right\|_2^2 \leq \|V_0\|_\infty \int_0^\infty x^{-\beta} e^{-2x} dx < \infty,$$

and so we have the result for $n = 0$. The case $n = -1$ follows by noting that $0 \leq T_{0,\delta}^{-1}(x) \leq \|V_0\|_\infty x$ and applying the same argument.

Now fix $n \geq 1$. We begin by noting that $\partial_x^n G_\delta(x, y) = \partial_y^n G_\delta^{r^n}(x, y)$, where $G^{r^n} = G$ if n is even and $G^{r^n} = G^r$ if n is odd. Splitting the range of integration and integrating by parts in the definition of $T_{0,\delta}$ gives

$$\begin{aligned} T_{0,\delta}^{(n)}(x) &= \sum_{i=0}^{n-1} (-1)^{n-1-i} \partial_y^{n-1-i} G_\delta^{r^n}(x, x/2) V_0^{(i)}(x/2) + \int_{x/2}^{\infty} G_\delta^{r^n}(x, y) V_0^{(n)}(y) dy \\ &\quad + \int_0^{x/2} \partial_x^n G_\delta(x, y) V_0(y) dy. \end{aligned} \quad (11)$$

We consider the weighted L^2 -norm of these three components.

Firstly by Lemma 4.6, there exists a numerical constant $D > 0$ such that

$$\begin{aligned} &\left| \partial_y^{n-1-i} G_\delta^{r^n}(x, x/2) V_0^{(i)}(x/2) w_{n-\beta/2}(x) \right| \\ &\leq D \sum_{\frac{n-1-i}{2} \leq j \leq n-1-i} x^{2j-n+1+i} \delta^{-j-1/2} e^{-\frac{x^2}{8\delta}} \left| V_0^{(i)}(x/2) \right| w_{i-\beta/2}(x) x^{n-i} \end{aligned}$$

therefore integrating over $x \in (0, \infty)$ and applying the Cauchy–Schwarz inequality

$$\begin{aligned} &\left\| \partial_y^{n-1-i} G_\delta^{r^n}(\cdot, \cdot/2) V_0^{(i)}(\cdot/2) w_{n-\beta/2} \right\|_2^2 \\ &\leq 2D \sum_{\frac{n-1-i}{2} \leq j \leq n-1-i} \int_0^{\infty} x^{4j+2} \delta^{-2j-1} e^{-\frac{x^2}{4\delta}} dx \left\| V_0^{(i)}(\cdot/2) w_{i-\beta/2} \right\|_2^2 \\ &\leq D' \sum_{\frac{n-1-i}{2} \leq j \leq n-1-i} \delta^{1/2} 2^{2i-\beta} \left\| V_0^{(i)} w_{i-\beta/2} \right\|_2^2 \rightarrow 0 \end{aligned} \quad (12)$$

as $\delta \rightarrow 0$, where $D' > 0$ is a further numerical constant.

For the second term we apply Cauchy–Schwarz to obtain

$$\begin{aligned} &\left\{ \int_{x/2}^{\infty} G_\delta^{r^n}(\cdot, y) V_0^{(n)}(y) dy \right\}^2 \\ &\leq \left\{ \int_0^{\infty} G_\delta^{r^n}(\cdot, y) w_{n-\beta/2}(y)^2 V_0^{(n)}(y)^2 dy \right\} \left\{ \int_{x/2}^{\infty} G_\delta^{r^n}(\cdot, y) y^{-2n+\beta} e^{2y} dy \right\} \\ &\leq 4 \cdot 2^{2n-\beta} x^{-2n+\beta} \left\{ \int_0^{\infty} p_\delta(x-y) w_{n-\beta/2}(y)^2 V_0^{(n)}(y)^2 dy \right\} \left\{ \int_{x/2}^{\infty} p_\delta(x-y) e^{2y} dy \right\}. \end{aligned}$$

For $\delta < 1$ we have $\int_{x/2}^{\infty} p_{\delta}(x-y) e^{2y} dy \leq D' e^{2x}$, for some numerical constant $D' > 0$, therefore

$$\begin{aligned} \left\| w_{n-\beta/2} \int_{\cdot/2}^{\infty} G_{\delta}^{r^n}(\cdot, y) V_0^{(n)}(y) dy \right\|_2^2 &\leq 2^{2n-\beta+2} D' \int_0^{\infty} \int_0^{\infty} p_{\delta}(x-y) dx w_{n-\beta/2}(y)^2 V_0^{(n)}(y)^2 dy \\ &\leq 2^{2n-\beta+2} D' \left\| w_{n-\beta/2} V_0^{(n)} \right\|_2^2. \end{aligned} \quad (13)$$

Finally using the fact that V_0 is bounded and Lemma 4.6, there exists a numerical constant $D'' > 0$ such that

$$\left| \int_0^{x/2} \partial_x^n G_{\delta}(x, y) V_0(y) dy \right| \leq D'' \|V_0\|_{\infty} x \sum_{n/2 \leq i \leq n} x^{2i-n} \delta^{-i-1/2} e^{-\frac{x^2}{8\delta}}$$

and therefore

$$\begin{aligned} \left\| w_{n-\beta/2} \int_0^{\cdot/2} \partial_x^n G_{\delta}(\cdot, y) V_0(y) dy \right\|_2^2 &\leq 2 (D'')^2 \|V_0\|_{\infty}^2 \sum_{n/2 \leq i \leq n} \int_0^{\infty} x^{4i-2n+2} \delta^{-2i-1} e^{-\frac{x^2}{8\delta}} dx \\ &\leq D''' \delta^{(1-\beta)/2} \rightarrow 0 \end{aligned} \quad (14)$$

as $\delta \rightarrow 0$, since $\beta < 1$. Here $D''' > 0$ is a further numerical constant.

Results (12), (13) and (14) give the required result by (11).

□

Lemma 4.8 For $n \geq 0$ and $\beta < \pi/\alpha - 1$

$$\liminf_{\delta \rightarrow 0} \mathbb{E} \left[\int_0^t \left\| w_{n-\beta/2} R_{s,\delta}^{(n+1)} \right\|_2^2 ds \right] = 0.$$

Proof: Interchanging differentiation and integration and applying Lemma 4.6 gives

$$\begin{aligned} \left| R_{s,\delta}^{(n+1)}(x) \right| &= \left| 2 \int_0^{\infty} \frac{\partial^{n+1} p_{\delta}}{\partial x^{n+1}}(x+y) V_s(y) dy \right| \\ &\leq D \sum_{\frac{n+1}{2} \leq i \leq n+1} \int_0^{\infty} (x+y)^{2i-n-1} \delta^{-i-1/2} e^{-\frac{x^2}{2\delta}} e^{-\frac{y^2}{2\delta}} V_s(y) dy, \end{aligned}$$

with $D > 0$ a numerical constant. Let $\eta \in (0, 1)$ and split the region of integration at $y = \delta^{\frac{1}{2}(1-\eta)}$, then for $\delta < 1$ we have $\delta^{-i} \leq \delta^{-n-1}$ and $e^{-\frac{y^2}{2\delta}} \leq e^{-\frac{y^2}{4}} e^{-1/4\delta\eta}$ on $y > \delta^{\frac{1}{2}(1-\eta)}$ so that

$$\begin{aligned} & \left| w_{n-\beta/2}(x) R_{s,\delta}^{(n+1)}(x) \right| \\ & \leq \left\{ D \sum_{\frac{n+1}{2} \leq i \leq n+1} \int_{\delta^{(1-\eta)/2}}^{\infty} (x+y)^{2i-n-1} e^{-\frac{x^2}{2}} e^{-\frac{y^2}{4}} \right\} \|V_0\|_{\infty} \delta^{-n-1/2} e^{-1/4\delta\eta} w_{n-\beta/2}(x) \quad (15) \\ & + D \sum_{\frac{n+1}{2} \leq i \leq n+1} \left(x + \delta^{(1-\eta)/2} \right)^{2i-n-1} \delta^{-i-1/2} \nu_s \left(0, \delta^{(1-\eta)/2} \right) w_{n-\beta/2}(x) e^{-\frac{x^2}{2\delta}}. \end{aligned}$$

The first term on the right-hand side of (15) vanishes as $\delta \rightarrow 0$, hence we have

$$\begin{aligned} \liminf_{\delta \rightarrow 0} \mathbb{E} \left[\int_0^t \left\| w_{n-\beta/2} R_{s,\delta}^{(n+1)} \right\|_2^2 ds \right] & \leq \liminf_{\delta \rightarrow 0} D' \sum_{\frac{n+1}{2} \leq i \leq n+1} \left\{ \int_0^{\infty} (x+1)^{4i-2n-2} x^{2n-\beta} e^{-x^2} dx \right. \\ & \quad \times \delta^{(1-\eta)(2i-n-1)} \delta^{-2i-1} \delta^{\frac{1}{2}(1-\eta)(3+\beta+\varepsilon)} \delta^{1/2} \delta^{n-\beta/2} \left. \right\} \quad (16) \end{aligned}$$

where we have used Theorem 2.6 with β replaced by $\beta + \varepsilon < \pi/\alpha - 1$ for small $\varepsilon > 0$, and $D' > 0$ is a numerical constant. (Note as we integrate over $s \in (0, t)$, the singular factor of $s^{-\gamma}$ from Theorem 2.6 integrates to a finite value because $\gamma \in (0, 1)$.) The δ term in (16) is bounded by

$$\delta^{\frac{1}{2}(1-\eta)\varepsilon - \eta(2n+5/2+\beta/2)}$$

for small δ . We choose η small enough so that

$$\frac{1}{2}(1-\eta)\varepsilon > \eta(2n+5/2+\beta/2)$$

then the limit on the right-hand side of (16) vanishes and gives the required result. □

4.2 The inductive proof for the regularity result

Initial case:

The initial case is when $n = -1$, and the result we shall demonstrate is:

$$\liminf_{\delta \rightarrow 0} \mathbb{E} \left[\int_0^T \left\| w_{-2-\beta/2} T_{t,\delta}^{(-1)} \right\|_2^2 dt \right] < \infty, \quad \text{for every } \beta < \pi/\alpha - 1. \quad (17)$$

By truncating the range of integration we have

$$T_{t,\delta}^{(-1)}(x) = \int_0^\infty \int_0^x G_\delta(z, y) dz V_t(y) dy = \nu_t(0, x) + I_{t,\delta}(x)$$

where $I_{t,\delta}(x) := \int_{2x}^\infty \int_0^x G_\delta(z, y) dz V_t(y) dy$. From Theorem 2.6 it is clear that

$$\mathbb{E} \left[\int_0^T \|w_{-2-\beta/2} \nu_t(0, \cdot)\|_2^2 dt \right] < \infty$$

and therefore we are done provided $\liminf_{\delta \rightarrow 0} \mathbb{E} \left[\int_0^T \|w_{-2-\beta/2} I_{t,\delta}\|_2^2 dt \right] < \infty$. Using the inequality

$$G_\delta(z, y) \leq \frac{2zy}{\sqrt{2\pi\delta^3}} \exp \left\{ -(z-y)^2 / 2\delta \right\},$$

we have a numerical constant $C > 0$ such that

$$I_{t,\delta}(x) \leq C \delta^{-3/2} x^2 e^{-x^2/2\delta} \int_0^\infty (y+2x) e^{-y^2/2\delta} V_t(y+2x) dy.$$

and so

$$\begin{aligned} & \|w_{-2-\beta/2} I_{t,\delta}\|_2^2 \\ & \leq \frac{C^2}{\delta^3} \int_0^\infty \int_0^\infty \int_0^\infty x^{-\beta} (y_1+2x) (y_2+2x) e^{-\frac{x^2+y_1^2+y_2^2}{2\delta}} V_t(y_1+2x) V_t(y_2+2x) dy_1 dy_2 dx. \end{aligned}$$

Now let $\eta \in (0, 1)$. Splitting the region of integration on $(0, \delta^{(1-\eta)/2})^3 \subseteq (0, \infty)^3$ and its complement, we see that on the complement the triple integral on the right-hand side above is bounded by a multiple of e^{-1/δ^η} , and so vanishes as $\delta \rightarrow 0$. Therefore we need only consider the region of integration $(0, \delta^{(1-\eta)/2})^3$, and so we have a numerical constant $C' > 0$ such that

$$\begin{aligned} \liminf_{\delta \rightarrow 0} \mathbb{E} \left[\int_0^T \|w_{-2-\beta/2} I_{t,\delta}\|_2^2 dt \right] & \leq C' \liminf_{\delta \rightarrow 0} \delta^{-(3+\beta)/2} \int_0^T \mathbb{E} \left[\nu_t \left(0, \delta^{(1-\eta)/2} \right)^2 \right] dt \\ & \leq B \liminf_{\delta \rightarrow 0} \delta^{(1-\eta)(3+\beta+\varepsilon)/2 - (3+\beta)/2} \end{aligned}$$

where the second line follows by Theorem 2.6 and $\varepsilon > 0$ is sufficiently small. Clearly we may choose $\eta > 0$ small enough so that the exponent of δ is positive, whereby the right-hand side vanishes and we have (17).

Inductive step:

Assume that for some $-1 \leq n \leq N-1$ we have

$$\liminf_{\delta \rightarrow 0} \mathbb{E} \left[\int_0^T \left\| w_{k-1-\beta/2} T_{t,\delta}^{(k)} \right\|_2^2 dt \right] < \infty, \quad \text{for every } \beta < \pi/\alpha - 1 \quad (18)$$

for $k = -1, 0, 1, \dots, n$. (We demonstrated the case $n = -1$ above.) The proof will be complete if we can show condition (18) holds for $k = n+1$, since then by induction the condition holds for all $k = 0, \dots, N$ and we can apply Lemma 4.4 to give the regularity result of Theorem 1.1 for the probabilistic solution ν .

Let us begin by multiplying throughout equation (9) by the square of the weighting function $w_{n-\beta/2}^2$ and integrating over $(0, \infty)$. To do so we apply Lemma 4.5 and arrive at

$$\begin{aligned} & \sigma_I^2 \mathbb{E} \left[\int_0^t \left\| w_{n-\beta/2} T_{s,\delta}^{(n+1)} \right\|_2^2 ds \right] \\ &= \left\| w_{n-\beta/2} T_{0,\delta}^{(n)} \right\|_2^2 - \mathbb{E} \left[\left\| w_{n-\beta/2} T_{t,\delta}^{(n)} \right\|_2^2 \right] \\ & - 2\mu \mathbb{E} \left[\int_0^t \left\| w_{n-\beta/2} T_{s,\delta}^{(n)} \right\|_2^2 ds \right] \\ & + 2\mu \left(n - \frac{\beta}{2} \right) \mathbb{E} \left[\int_0^t \left\| w_{n-1/2-\beta/2} T_{s,\delta}^{(n)} \right\|_2^2 ds \right] \\ & + 2(\sigma_M^2 + \sigma_I^2) \mathbb{E} \left[\int_0^t \left\| w_{n-\beta/2} T_{s,\delta}^{(n)} \right\|_2^2 ds \right] \\ & - 4(\sigma_M^2 + \sigma_I^2) \left(n - \frac{\beta}{2} \right) \mathbb{E} \left[\int_0^t \left\| w_{n-1/2-\beta/2} T_{s,\delta}^{(n)} \right\|_2^2 ds \right] \\ & + 2(\sigma_M^2 + \sigma_I^2) \left(n - \frac{\beta}{2} \right) \left(n - \frac{\beta+1}{2} \right) \mathbb{E} \left[\int_0^t \left\| w_{n-1-\beta/2} T_{s,\delta}^{(n)} \right\|_2^2 ds \right] \\ & + \sigma_M^2 \mathbb{E} \left[\int_0^t \left\| w_{n-\beta/2} R_{s,\delta}^{(n+1)} \right\|_2^2 ds \right] \\ & + 2\mu \mathbb{E} \left[\int_0^t \int_0^\infty w_{n-\beta/2}^2 T_{s,\delta}^{(n)} R_{s,\delta}^{(n+1)} ds \right] \end{aligned} \quad (19)$$

Eliminating the negative terms from the right-hand side and applying the Cauchy–Schwarz inequality

gives

$$\begin{aligned}
& \sigma_I^2 \mathbb{E} \left[\int_0^t \left\| w_{n-\beta/2} T_{s,\delta}^{(n+1)} \right\|_2^2 ds \right] \\
& \leq \left\| w_{n-\beta/2} T_{0,\delta}^{(n)} \right\|_2^2 \\
& - 2\mu \mathbb{E} \left[\int_0^t \left\| w_{n-\beta/2} T_{s,\delta}^{(n)} \right\|_2^2 ds \right] \\
& + 2\mu \left(n - \frac{\beta}{2} \right) \mathbb{E} \left[\int_0^t \left\| w_{n-1/2-\beta/2} T_{s,\delta}^{(n)} \right\|_2^2 ds \right] \\
& + 2 \left(\sigma_M^2 + \sigma_I^2 \right) \mathbb{E} \left[\int_0^t \left\| w_{n-\beta/2} T_{s,\delta}^{(n)} \right\|_2^2 ds \right] \\
& + 2 \left(\sigma_M^2 + \sigma_I^2 \right) \left(n - \frac{\beta}{2} \right) \left(n - \frac{\beta+1}{2} \right) \mathbb{E} \left[\int_0^t \left\| w_{n-1-\beta/2} T_{s,\delta}^{(n)} \right\|_2^2 ds \right] \\
& + \sigma_M^2 \mathbb{E} \left[\int_0^t \left\| w_{n-\beta/2} R_{s,\delta}^{(n+1)} \right\|_2^2 ds \right] \\
& + 2 |\mu| \left\{ \mathbb{E} \left[\int_0^t \left\| w_{n-\beta/2} T_{s,\delta}^{(n)} \right\|_2^2 ds \right] \right\}^{1/2} \left\{ \mathbb{E} \left[\int_0^t \left\| w_{n-\beta/2} R_{s,\delta}^{(n+1)} \right\|_2^2 ds \right] \right\}^{1/2}.
\end{aligned} \tag{20}$$

We are done provided the seven terms on the right-hand side of inequality (20) have a finite limit-infimum as $\delta \downarrow 0$. This is true of the second to fifth terms by the inductive hypothesis (18), and with this the final two terms vanish by Lemma 4.8. From Lemma 4.7 the first term remains finite in the limit-infimum, hence we have shown the probabilistic solution ν satisfies the regularity properties in Theorem 1.1.

4.3 Uniqueness

It remains to show that the probabilistic solution is indeed the unique finite-measure valued solution of (1). Suppose we have another finite-measure valued solution $\bar{\nu}$. Define the signed-measure valued process u as the difference between the solutions:

$$u_t(S) := \nu_t(S) - \bar{\nu}_t(S).$$

By linearity of (1), provided that the technical lemmas in Section 5.1 hold if we replace ν by u , then we arrive at (19) and the same limit results hold. Whence in the case $n = 0$ and $\beta = 0$ we have

$$\sigma_I^2 \mathbb{E} \left[\int_0^t \left\| U'_{s,\delta} \right\|_2^2 ds \right] = -\mathbb{E} \left[\left\| U_{t,\delta} \right\|_2^2 \right] + 2 \left(\sigma_M^2 + \sigma_I^2 - \mu \right) \int_0^t \mathbb{E} \left[\left\| U_{s,\delta} \right\|_2^2 \right] ds + f(t, \delta), \tag{21}$$

where $U_{t,\delta}$ denotes the function obtained by applying the kernel smoothing operator to u_t and $f(t, \delta) \rightarrow 0$ as $\delta \rightarrow 0$ uniformly in $t \in [0, T]$. Observe that the initial condition vanishes because

$u_0 = 0$. Since the left-hand side of (21) is non-negative, we have

$$\liminf_{\delta \rightarrow 0} \mathbb{E} \left[\|U_{t,\delta}\|_2^2 \right] = 0$$

for all $t \in [0, T]$ by Gronwall's Lemma. This is then enough to show $u \equiv 0$ provided Lemma 4.4 is applicable for u .

It therefore remains to show the working in Section 4.1 is valid for ν replaced by u and the specific choice $N = 0$, $\beta = 0$. Firstly observe that by taking expectation over (1), the (deterministic) measure-valued process $(\mathbb{E}[u_t])_{t \in [0, T]}$ satisfies the (weak) heat equation in the half-line with zero boundary condition and zero initial condition. As u_t is a finite-measure, the uniqueness of solutions to the heat equation forces $\mathbb{E}[u_t] \equiv 0$, and so $\mathbb{E}[\nu_t] \equiv \mathbb{E}[\bar{\nu}_t]$. Likewise, by considering the two-dimensional heat equation (with a correlation term), we conclude that $\mathbb{E}[\nu_t(S_1)\nu_t(S_2)] = \mathbb{E}[\bar{\nu}_t(S_1)\bar{\nu}_t(S_2)]$, for all measurable $S_1, S_2 \subseteq (0, \infty)$. Therefore by the Hahn Decomposition Theorem we can write

$$\mathbb{E} \left[u_t(0, \varepsilon)^2 \right] \leq \mathbb{E} \left[\nu_t(0, \varepsilon)^2 \right] + \mathbb{E} \left[\bar{\nu}_t(0, \varepsilon)^2 \right] = 2\mathbb{E} \left[\nu_t(0, \varepsilon)^2 \right].$$

This is enough to guarantee Lemma 4.8 holds for u . As Lemma 4.3 may no longer be true for u , the remaining facts that need to be checked are whether the limit $\lim_{x \downarrow 0} U_{t,\delta}(x) = 0$ from Lemma 4.4 and the limits

$$\begin{aligned} \lim_{x \downarrow 0} U_{t,\delta}(x) w_{-1/2}(x) &= 0 \\ \lim_{x \downarrow 0} U_{t,\delta}(x) U'_{t,\delta}(x) &= 0 \end{aligned}$$

from Lemma 4.5 remain valid. However it is easy to see that

$$\begin{aligned} |U_{t,\delta}(x)| &\leq \int_0^\infty \{p_\delta(x-y) - p_\delta(x+y)\} |u_t|(dy) \\ &= \frac{1}{\sqrt{2\pi\delta}} \int_0^\infty \exp\left\{-\frac{(x-y)^2}{2\delta}\right\} \left(1 - \exp\left\{-\frac{2xy}{\delta}\right\}\right) |u_t|(dy) \\ &\leq \frac{2x}{\delta\sqrt{2\pi\delta}} \int_0^\infty \exp\left\{-\frac{(x-y)^2}{2\delta}\right\} y |u_t|(dy) \end{aligned}$$

where $|u_t| \leq \nu_t + \bar{\nu}_t$. For $x < 1$ the integrand is bounded and hence the integral is finite since $|u_t|$ is a finite measure. We therefore conclude that $U_{t,\delta}(x) = \mathcal{O}(x)$ as $x \rightarrow 0$, which is enough to complete the proof.

□

5 Proof of Theorem 1.2

This theorem relies on the fact that the estimate in Theorem 2.6 is sharp. Specifically, Lemma 5.1 of [6] states that if V_0 is supported on a finite interval away from zero, then for every $\eta > 0$ there exist $C(\eta), \varepsilon_0(\eta) > 0$ such that for all $0 < \varepsilon < \varepsilon_0(\eta)$ and $t \in [0, T]$

$$\mathbb{E} \left[\nu_t(0, \varepsilon)^2 \right] \geq C(\eta) \varepsilon^{2+\pi/\alpha+\eta}. \quad (22)$$

It is clear that if $\tilde{V}_0 \leq V_0$ then the empirical measure $\tilde{\nu}$ arising from starting the system from $\tilde{V}_0$ satisfies $\tilde{\nu}_t \leq \nu_t$. Since we can certainly find a constant $a > 0$ such that $\tilde{V}_0 := V_0 \mathbf{1}_{(a, \infty)}$ is non-zero, we see that (22) remains valid if we drop the assumption on the support of V_0 .

For a contradiction, suppose that we have $\beta > \pi/\alpha - 1$ and a $k \in \{0, 1, \dots, N\}$ for which

$$\mathbb{E} \left[\int_0^T \left\| w_{k-1-\beta/2} V_t^{(k)} \right\|_2^2 dt \right] < \infty.$$

Then in particular

$$\mathbb{E} \left[\int_0^T \int_0^1 x^{2(k-1)-\tilde{\beta}} V_t^{(k)}(x)^2 dx dt \right] < \infty$$

for any $\tilde{\beta} < \beta$, so let us fix $\tilde{\beta} \in (\pi/\alpha - 1, 1)$. Also take $\varepsilon > 0$ small enough so that we have $\tilde{\beta} - \varepsilon \in (\pi/\alpha - 1, 1)$. Then for $k \geq 2$

$$\int_0^1 x^{2(k-2)-\tilde{\beta}+\varepsilon} dx < \infty$$

and since

$$\begin{aligned} \left(V_t^{(k-1)}(x) - V_t^{(k-1)}(1) \right)^2 &= \left(\int_x^1 V_t^{(k)}(y) dy \right)^2 \\ &\leq \int_0^1 x^{2(k-1)-\tilde{\beta}} V_t^{(k)}(x)^2 dx \cdot \int_x^1 y^{-2k+2+\tilde{\beta}} dy \\ &\leq \int_0^1 x^{2(k-1)-\tilde{\beta}} V_t^{(k)}(x)^2 dx \cdot \frac{x^{-2k+3+\tilde{\beta}}}{2k-3-\tilde{\beta}} \end{aligned} \quad (23)$$

applying the triangle inequality, multiplying by $x^{2k-4-\tilde{\beta}+\varepsilon}$ and integrating over $\Omega \times [0, T] \times (0, 1)$ gives

$$\mathbb{E} \left[\int_0^T \int_0^1 x^{2(k-2)-\tilde{\beta}+\varepsilon} V_t^{(k-1)}(x)^2 dx dt \right] < \infty.$$

Repeating this argument eventually yields

$$\mathbb{E} \left[\int_0^T \int_0^1 x^{-\tilde{\beta}+\varepsilon} V_t'(x)^2 dx dt \right] < \infty$$

for all allowable ε . If we return to (23) with $k = 1$ and take the upper limit of the integral to be x and the lower limit 0, then since $V_t(0) = 0$ we arrive at

$$\mathbb{E} \left[\int_0^T \int_0^1 x^{-2-\tilde{\beta}+\varepsilon} V_t(x)^2 dx dt \right] < \infty.$$

Repeating this manipulation a final time gives the following for $x < 1$:

$$\begin{aligned} \mathbb{E} \left[\int_0^T \nu_t(0, x)^2 dt \right] &\leq \mathbb{E} \left[\int_0^T \int_0^1 x^{-2-\tilde{\beta}+\varepsilon} V_t(x)^2 dx dt \right] \int_0^x y^{2+\tilde{\beta}-\varepsilon} dy \\ &= D(\tilde{\beta}, \varepsilon) x^{3+\tilde{\beta}-\varepsilon} \end{aligned}$$

with $D(\tilde{\beta}, \varepsilon) \in (0, \infty)$ a constant. This is a contradiction of (22) for the case $\eta = (1 - \pi/\alpha) + (\tilde{\beta} - \varepsilon) > 0$. Hence we have Theorem 1.2.

□

References

- [1] Bain, A. and Crisan, D. (2009) *Fundamentals of stochastic filtering*, Springer.
- [2] Bertoin, J. (1996) *L $\tilde{A}$ @vy Processes*, Cambridge University Press.
- [3] Bichteler, K., Lin, S.J. (1995) On the stochastic Fubini theorem, *Stochastics Rep.* 54(3-4): 271-279.
- [4] Bush, N., Hambly, B.M., Haworth, H., Jin, L., Reisinger, C. (2011) Stochastic evolution equations in portfolio credit modelling with applications to exotic credit products, *SIAM J. Fin. Math.* 2, 627–664.
- [5] Iyengar, S. (1985) Hitting lines with two-dimensional Brownian motion, *SIAM J. Appl. Math.*, 45, 983-989.
- [6] Jin, L. (2010) Particle systems and SPDEs with application to credit modelling. *D.Phil. Thesis*, University of Oxford.
- [7] Kim, K-H., Krylov, N. V. (2004) On SPDEs with Variable Coefficients in One Space Dimension, *Potential Analysis*, 21: 209-239.
- [8] Krylov, N. V. (1994) A W_2^n -theory of the Dirichlet problem for SPDEs in general smooth domains, *Probab. Theory Relat. Fields*, 389-421.
- [9] Krylov, N. V. (2003) Brownian Trajectory is a Regular Lateral Boundary for the Heat Equation, *SIAM J. Math. Anal.*, Vol. 34, 1167-1182.
- [10] Krylov, N. V., Lototsky, S. V. (1998) A Sobolev Space Theory of SPDEs with Constant Coefficients on a Half Line, *SIAM J. Math. Anal.*, Vol. 30, 298-325.
- [11] Kurtz, T. G., Xiong, J. (1999) Particle representations for a class of non-linear SPDEs, *Stoch. Proc. Appl.*, 83, 103-126.
- [12] Metzler, A. (2010) On the first passage problem for correlated Brownian motion, *Statistics and Probability Letters*, 80, 277-284.